

INSTRUCTIONS FOR USE

OLPRUVA™ (ol proo vah)

(sodium phenylbutyrate)




for oral suspension

Read this Instructions for Use before taking OLPRUVA oral suspension and each time you get a refill. There may be new information. This Instructions for Use does not take the place of talking to your healthcare provider about your medical condition or treatment. Talk to your healthcare provider or pharmacist if you have any questions about how to take a dose of OLPRUVA.

This Instructions for Use contains information on how to prepare and take 1 dose of OLPRUVA.

Supplies needed to take 1 dose of OLPRUVA as prescribed:

- One dosage envelope containing OLPRUVA and Mix-Aid packets. The contents of 1 envelope equals 1 full dose.
- An open drinking cup
- A spoon
- Water

Step 1: Get an open drinking cup and spoon (See Figure A).	 Figure A
Step 2: Add about 4 ounces of water to an open drinking cup (See Figure B).	 Figure B
Step 3: Remove 1 dosage envelope from the kit (See Figure C). Note: Each kit is divided into 30 individual sections. Each section contains 3 dosage envelopes for a total of 90 dosage envelopes.	 Figure C

Step 4: Open the dosage envelope and remove all packets (1 packet of Mix-Aid and 1 or 2 packets of OLPRUVA) (See **Figure D**).

Note: All packets in the dosage envelope must be used for 1 full dose.




Figure D

Step 5: To open the Mix-Aid packet, tear or cut with scissors, straight across from the notch (See **Figures E and F**).



Figure E



Figure F

Step 6: Add the full contents of the Mix-Aid packet to the water (in the open drinking cup) and stir with the spoon for at least 30 seconds (See **Figure G**).

Note: The contents will not dissolve but will make the water thicker so that OLPRUVA does not sink to the bottom of the cup.

Do not drink yet.



Figure G




Step 7: To open the OLPRUVA packet(s), fold at the notch and tear or cut with scissors. Add the full contents of the OLPRUVA packet(s) to the open drinking cup containing the mixture from Step 6 (the water and Mix-Aid) and stir for 15 seconds (See **Figure H**).

Drink the entire contents of the open drinking cup within 5 minutes to help prevent the coating from dissolving (See **Figure I**).

Note: The entire contents of each packet must be used for 1 full dose.



Figure H

	 <p>Figure I</p>
<p>Step 8: To make sure that you get the full dose, pour another 4 ounces of water in the open drinking cup, stir (See Figure J) and drink the entire contents (See Figure K).</p> <p>OLPRUVA suspension is taken by mouth only. Do not take or give OLPRUVA suspension in a gastrostomy or nasogastric tube.</p> <p>The mixed OLPRUVA suspension should be thrown away (discarded) after 30 minutes, if not used right away.</p>	 <p>Figure J</p>  <p>Figure K</p>

How should I store OLPRUVA?

- Store OLPRUVA at room temperature between 68°F and 77°F (20°C and 25°C).
- **Keep OLPRUVA and all medicines out of the reach of children.**



acertherapeutics

Manufactured for:

Acer Therapeutics Inc.
300 Washington St.
Newton, MA 02458

For more information, go to www.OLPRUVA.com or call Acer Therapeutics Inc. at 1-844-600-2237.

© 2022 Acer Therapeutics Inc. All rights reserved.